

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 December 2001 (27.12.2001)

PCT

(10) International Publication Number
WO 01/98259 A1

- (51) International Patent Classification⁷: C07C 237/26, A61K 31/165, A61P 31/04
- (21) International Application Number: PCT/US00/16632
- (22) International Filing Date: 16 June 2000 (16.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (71) Applicant: **TRUSTEES OF TUFTS COLLEGE**
[US/US]; Ballou Hall, 4th Floor, Medford, MA 02155 (US).
- (72) Inventors: **NELSON, Mark**; 735 Worcester Road, Wellesley, MA 02481 (US). **RENNIE, Glen**; 1592 Beacon Street, Apt. 3, Brookline, MA 02146 (US).
- (74) Agents: **HANLEY, Elizabeth, A. et al.**; Lahive & Cockfield, LLP, 28 State Street, Boston, MA 02109 (US).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/98259 A1

(54) Title: 7-PHENYL-SUBSTITUTED TETRACYCLINE COMPOUNDS

(57) Abstract: 7-phenyl-substituted tetracycline compounds, methods of treating tetracycline responsive states, and pharmaceutical compositions containing the 7-phenyl-substituted tetracycline compounds are described.

7-PHENYL-SUBSTITUTED TETRACYCLINE COMPOUNDS

BACKGROUND OF THE INVENTION


The development of the tetracycline antibiotics was the direct result of a systematic
5 screening of soil specimens collected from many parts of the world for evidence of
microorganisms capable of producing bacteriocidal and/or bacteriostatic compositions. The
first of these novel compounds was introduced in 1948 under the name chlortetracycline. Two
years later, oxytetracycline became available. The elucidation of the chemical structure of
these compounds confirmed their similarity and furnished the analytical basis for the
10 production of a third member of this group in 1952, tetracycline. A new family of tetracycline
compounds, without the ring-attached methyl group present in earlier tetracyclines, was
prepared in 1957 and became publicly available in 1967; and minocycline was in use by 1972.

Recently, research efforts have focused on developing new tetracycline antibiotic
compositions effective under varying therapeutic conditions and routes of administration. New
15 tetracycline analogues have also been investigated which may prove to be equal to or more
effective than the originally introduced tetracycline compounds. Examples include U.S. Patent
Nos. 3,957,980; 3,674,859; 2,980,584; 2,990,331; 3,062,717; 3,557,280; 4,018,889; 4,024,272;
4,126,680; 3,454,697; and 3,165,531. These patents are representative of the range of
pharmaceutically active tetracycline and tetracycline analogue compositions.

20 Historically, soon after their initial development and introduction, the tetracyclines were
found to be highly effective pharmacologically against rickettsiae; a number of gram-positive
and gram-negative bacteria; and the agents responsible for lymphogranuloma venereum,
inclusion conjunctivitis, and psittacosis. Hence, tetracyclines became known as "broad
spectrum" antibiotics. With the subsequent establishment of their in vitro antimicrobial
25 activity, effectiveness in experimental infections, and pharmacological properties, the
tetracyclines as a class rapidly became widely used for therapeutic purposes. However, this
widespread use of tetracyclines for both major and minor illnesses and diseases led directly to
the emergence of resistance to these antibiotics even among highly susceptible bacterial species
both commensal and pathogenic (e.g., pneumococci and Salmonella). The rise of tetracycline-
30 resistant organisms has resulted in a general decline in use of tetracyclines and tetracycline
analogue compositions as antibiotics of choice.

SUMMARY OF THE INVENTION:

The invention pertains to 7-substituted tetracycline compounds of the formula:



(I)

5 wherein:

R^4 and $R^{4'}$ are each alkyl;

R^5 is hydrogen, hydroxyl, or a prodrug moiety;


R^6 and $R^{6'}$ are each independently hydrogen, hydroxyl, alkyl, or taken together, alkenyl;

R^7 is halo substituted or unsubstituted phenyl; and pharmaceutically acceptable salts thereof. In another embodiment, R^7 is mono-, di-, or tri- halo substituted phenyl. In another embodiment, R^7 is 2- halo substituted phenyl. In yet another embodiment, R^7 is 3- halo substituted phenyl. In a further embodiment, R^7 is 4- halo substituted phenyl.

The invention also pertains to a method for treating a tetracycline responsive state in a mammal, by administering to a mammal a compound of formula I. In another aspect, the invention relates to the use of a compound of formula I to treat a tetracycline responsive state. The invention also pertains to pharmaceutical compositions comprising a compound of formula I, and to the use of a compound of formula I in the manufacture of a medicament to treat a tetracycline responsive state.

DETAILED DESCRIPTION OF THE INVENTION:

20 The invention pertains to 7-substituted tetracycline compounds of the formula:



(I)

wherein:

R^4 and $R^{4'}$ are each alkyl;








R^5 is hydrogen, hydroxyl, or a prodrug moiety;

R^6 and $R^{6'}$ are each independently hydrogen, hydroxyl, alkyl, or taken together, alkenyl;

R^7 is halo substituted or unsubstituted phenyl; and pharmaceutically acceptable salts thereof. In another embodiment, R^7 is mono-, di-, or tri- halo substituted phenyl. In another embodiment, R^7 is 2- halo substituted phenyl. In yet another embodiment, R^7 is 3- halo substituted phenyl. In a further embodiment, R^7 is 4- halo substituted phenyl.

The term "tetracycline compound" includes compounds with a similar ring structure to tetracycline, such as those included in formula I. Some examples of tetracycline compounds which can be modified to include a substituent at position 7 include tetracycline, oxytetracycline, demeclocycline, methacycline, sancycline, and doxycycline; however, other derivatives and analogues comprising a similar ring structure are also included. Table 1 depicts tetracycline and several known tetracycline derivatives.

TABLE I

 <p style="text-align: center;">Tetracycline</p>	
 <p style="text-align: center;">Oxytetracycline</p>	 <p style="text-align: center;">Demeclocycline</p>
 <p style="text-align: center;">Methacycline</p>	 <p style="text-align: center;">Doxycycline</p>
 <p style="text-align: center;">Chlorotetracycline</p>	 <p style="text-align: center;">Minocycline</p>

The term "7-substituted tetracycline compounds" includes tetracycline compounds with a phenyl substituent at the 7 position. In an embodiment, the substituted tetracycline

5 compound is substituted tetracycline (e.g., wherein R^4 and $R^{4'}$ are methyl, R^5 is hydrogen, R^6 is methyl and $R^{6'}$ is hydroxyl); substituted doxycycline (e.g., wherein R^4 and $R^{4'}$ are methyl, R^5 is hydroxyl, R^6 is methyl and $R^{6'}$ is hydrogen); or substituted sancycline (wherein R^4 and $R^{4'}$ are methyl; R^5 is hydrogen and R^6 and $R^{6'}$ are hydrogen atoms). In another embodiment, the compound is a derivative of tetracycline, sancycline, doxycycline, oxytetracycline, or

10 methacycline. In one embodiment, R^5 , R^6 and $R^{6'}$ are each hydrogen and R^4 and $R^{4'}$ are each methyl.

In yet another further embodiment, R⁷ is unsubstituted phenyl. Examples of tetracycline compounds with this R⁷ substituent include 7-phenyl sancycline and 7,9 diphenyl sancycline.

5 In yet another embodiment, R⁷ is halo substituted phenyl. The halo substituent can be, for example, chlorine, fluorine, bromine, or iodine, as well as mono-, di- or tri- halo substituted lower alkyl group, e.g., mono-, di- or tri- halo substituted methyl. In certain embodiments, the halo substitution of the phenyl substituent enhances the ability of the tetracycline compound to perform its intended function, e.g., treat tetracycline responsive states.

10 In a further embodiment, R⁷ is mono-, di-, or tri- halo substituted phenyl. In certain embodiments, the 7-substituted tetracycline compound is 7-(2,4-difluorophenyl) sancycline, 7-(2,4-dichlorophenyl) sancycline, 7-(2,4-dibromophenyl) sancycline, or 7-(2,4-diiodophenyl) sancycline. In other embodiments, the 7-substituted tetracycline compound is 7-(2,6-difluorophenyl) sancycline, 7-(2,6-dichlorophenyl) sancycline, 7-(2,6-dibromophenyl) sancycline, or 7-(2,6-diiodophenyl) sancycline.

15 In another embodiment, R⁷ is 2-halo substituted phenyl, e.g., 7-(2-fluorophenyl) sancycline, 7-(2-chlorophenyl) sancycline, 7-(2-bromophenyl) sancycline, or 7-(2-iodophenyl) sancycline.


In yet another embodiment, R⁷ is 3-halo substituted phenyl, e.g., 7-(3-fluorophenyl) sancycline, 7-(3-chlorophenyl) sancycline, 7-(3-bromophenyl) sancycline, or 7-(3-iodophenyl) sancycline.
20

In further embodiment, R⁷ is 4-halo substituted phenyl, e.g., 7-(4-fluorophenyl) sancycline, 7-(4-chlorophenyl) sancycline, 7-(4-bromophenyl) sancycline, or 7-(4-iodophenyl) sancycline.

25 In yet another embodiment, R⁷ is a mono-, di-, or tri- substituted phenyl where the substituent is a mono-, di- or tri- halo substituted lower alkyl group, e.g., mono-, di- or tri- halo substituted methyl. For example, the compound may be 7-(4-trichloromethylphenyl) sancycline, 7-(4-trifluoromethylphenyl) sancycline, 7-(4-tribromomethylphenyl) sancycline, or 7-(4-triiodomethylphenyl) sancycline.

30 The 7-substituted compounds of the invention can be synthesized by methods known in the art and/or as described herein. In Scheme 1, a general synthetic scheme is outlined using a Suzuki coupling of a boronic acid with an iodo tetracycline compound. Although the reaction

is shown for sancycline, a similar procedure can be used for other tetracycline compounds. Furthermore, other aryl coupling reactions known in the art may also be used.



SCHEME 1

5 As shown in Scheme 1, an iodosancycline compound can be synthesized from unsubstituted sancycline by treating it with at least one equivalent N-iodosuccinimide (NIS) under acidic conditions. The reaction is then quenched, and the resulting 7-iodosancycline can then be purified using standard techniques known in the art. The 7-iodosancycline can then be further reacted with a boronic acid, as shown in Scheme 1. 7-iodosancycline, a palladium catalyst (such as Pd(OAc)₂), is dissolved in a solvent and treated with aqueous sodium carbonate, and the boronic acid. The resulting compound can then be purified using techniques known in the art such as preparative HPLC and characterized.

The compounds of the invention can also be synthesized using Stille cross couplings. Stille cross couplings can be performed using an appropriate tin reagent (e.g., R-SnBu₃) and a halogenated tetracycline compound, (e.g., 7-iodosancycline). The tin reagent and the iodotetracycline compound can be treated with a palladium catalyst (e.g., Pd(PPh₃)₂Cl₂ or Pd(AsPh₃)₂Cl₂) and, optionally, with an additional copper salt, e.g., CuI. The resulting compound can then be purified using techniques known in the art. The synthesis of the compounds of the invention are described in more detail in Example 1. . Synthetic methods also are described in related U.S. provisional application entitled "7-Phenyl-Substituted Tetracycline Compounds", filed on even date herewith, the contents of which are expressly incorporated by reference.

The term "alkyl" includes saturated aliphatic groups, including straight-chain alkyl groups (e.g., methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, nonyl, decyl, etc.), branched-chain alkyl groups (isopropyl, tert-butyl, isobutyl, etc.), cycloalkyl (alicyclic) groups (cyclopropyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl), alkyl substituted cycloalkyl groups, and cycloalkyl substituted alkyl groups. The term alkyl further includes alkyl groups, which comprise oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more carbons of the hydrocarbon backbone. In certain embodiments, a straight chain or branched chain alkyl has 6 or fewer carbon atoms in its backbone (e.g., C₁-C₆ for straight chain, C₃-C₆ for branched chain), and more preferably 4 or fewer. Likewise, preferred cycloalkyls have from 3-8 carbon atoms in their ring structure, and more preferably have 5 or 6 carbons in the ring structure. The term C₁-C₆ includes alkyl groups containing 1 to 6 carbon atoms.

Moreover, the term alkyl includes both "unsubstituted alkyls" and "substituted alkyls", the latter of which refers to alkyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl, alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonate, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Cycloalkyls can be further substituted, e.g., with the substituents described above. An "alkylaryl" or an "aralkyl" moiety is an alkyl substituted with an aryl (e.g., phenylmethyl (benzyl)). The term "alkyl" also includes the side chains of natural and unnatural amino acids.

The term "aryl" includes groups with aromaticity, including 5- and 6-membered single-ring aromatic groups that may include from zero to four heteroatoms as well as multicyclic systems with at least one aromatic ring. Examples of aryl groups include benzene, phenyl, pyrrole, furan, thiophene, thiazole, isothiazole, imidazole, triazole, tetrazole, pyrazole, oxazole, isooxazole, pyridine, pyrazine, pyridazine, and pyrimidine, and the like. Furthermore, the term "aryl" includes multicyclic aryl groups, e.g., tricyclic, bicyclic, e.g., naphthalene, benzoxazole,

benzodioxazole, benzothiazole, benzoimidazole, benzothiophene, methylenedioxyphenyl, quinoline, isoquinoline, naphthridine, indole, benzofuran, purine, benzofuran, deazapurine, or indolizine. Those aryl groups having heteroatoms in the ring structure may also be referred to as "aryl heterocycles", "heterocycles," "heteroaryls" or "heteroaromatics". The aromatic ring
5 can be substituted at one or more ring positions with such substituents as described above, as for example, halogen, hydroxyl, alkoxy, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy, carboxylate, alkylcarbonyl, alkylaminocarbonyl, aralkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, alkenylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylthiocarbonyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety. Aryl groups can also be fused or bridged with
15 alicyclic or heterocyclic rings which are not aromatic so as to form a multicyclic system (e.g., tetralin, methylenedioxyphenyl).

The term "alkenyl" includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but that contain at least one double bond.

For example, the term "alkenyl" includes straight-chain alkenyl groups (e.g., ethenyl, propenyl, butenyl, pentenyl, hexenyl, heptenyl, octenyl, nonenyl, decenyl, etc.), branched-chain
20 alkenyl groups, cycloalkenyl (alicyclic) groups (cyclopropenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl), alkyl or alkenyl substituted cycloalkenyl groups, and cycloalkyl or cycloalkenyl substituted alkenyl groups. The term alkenyl further includes alkenyl groups which include oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more carbons of
25 the hydrocarbon backbone. In certain embodiments, a straight chain or branched chain alkenyl group has 6 or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). Likewise, cycloalkenyl groups may have from 3-8 carbon atoms in their ring structure, and more preferably have 5 or 6 carbons in the ring structure. The term C₂-C₆ includes alkenyl groups containing 2 to 6 carbon atoms.

Moreover, the term alkenyl includes both "unsubstituted alkenyls" and "substituted alkenyls", the latter of which refers to alkenyl moieties having substituents replacing a
30 hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include,

for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including
5 alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

10 The term "alkynyl" includes unsaturated aliphatic groups analogous in length and possible substitution to the alkyls described above, but which contain at least one triple bond.

For example, the term "alkynyl" includes straight-chain alkynyl groups (e.g., ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl, octynyl, nonynyl, decynyl, etc.), branched-chain alkynyl groups, and cycloalkyl or cycloalkenyl substituted alkynyl groups. The term
15 alkynyl further includes alkynyl groups which include oxygen, nitrogen, sulfur or phosphorous atoms replacing one or more carbons of the hydrocarbon backbone. In certain embodiments, a straight chain or branched chain alkynyl group has 6 or fewer carbon atoms in its backbone (e.g., C₂-C₆ for straight chain, C₃-C₆ for branched chain). The term C₂-C₆ includes alkynyl groups containing 2 to 6 carbon atoms.

20 Moreover, the term alkynyl includes both "unsubstituted alkynyls" and "substituted alkynyls", the latter of which refers to alkynyl moieties having substituents replacing a hydrogen on one or more carbons of the hydrocarbon backbone. Such substituents can include, for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl,
25 arylcarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido,
30 nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

Unless the number of carbons is otherwise specified, "lower alkyl" as used herein means an alkyl group, as defined above, but having from one to five carbon atoms in its backbone structure. "Lower alkenyl" and "lower alkynyl" have chain lengths of, for example, 2-5 carbon atoms.

5 The term "acyl" includes compounds and moieties which contain the acyl radical ($\text{CH}_3\text{CO}-$) or a carbonyl group. The term "substituted acyl" includes acyl groups where one or more of the hydrogen atoms are replaced by for example, alkyl groups, alkynyl groups, halogens, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, 10 alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl, alkoxy, phosphate, phosphonate, phosphinate, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonate, sulfamoyl, sulfonamido, nitro, 15 trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moiety.

The term "acylamino" includes moieties wherein an acyl moiety is bonded to an amino group. For example, the term includes alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido groups.

20 The term "aroyl" includes compounds and moieties with an aryl or heteroaromatic moiety bound to a carbonyl group. Examples of aroyl groups include phenylcarboxy, naphthyl carboxy, etc.

The terms "alkoxyalkyl", "alkylaminoalkyl" and "thioalkoxyalkyl" include alkyl groups, as described above, which further include oxygen, nitrogen or sulfur atoms replacing one or more carbons of the hydrocarbon backbone, e.g., oxygen, nitrogen or sulfur atoms.

25 The term "alkoxy" includes substituted and unsubstituted alkyl, alkenyl, and alkynyl groups covalently linked to an oxygen atom. Examples of alkoxy groups include methoxy, ethoxy, isopropoxy, propoxy, butoxy, and pentoxy groups. Examples of substituted alkoxy groups include halogenated alkoxy groups. The alkoxy groups can be substituted with groups such as alkenyl, alkynyl, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, 30 alkoxy carbonyloxy, aryloxy carbonyloxy, carboxylate, alkylcarbonyl, arylcarbonyl, alkoxy carbonyl, aminocarbonyl, alkylaminocarbonyl, dialkylaminocarbonyl, alkylthiocarbonyl,

alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including alkylamino, dialkylamino, arylamino, diarylamino, and alkylarylamino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl, alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, 5 nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkylaryl, or an aromatic or heteroaromatic moieties. Examples of halogen substituted alkoxy groups include, but are not limited to, fluoromethoxy, difluoromethoxy, trifluoromethoxy, chloromethoxy, dichloromethoxy, trichloromethoxy, etc.

The term "amine" or "amino" includes compounds where a nitrogen atom is covalently 10 bonded to at least one carbon or heteroatom. The term "alkylamino" includes groups and compounds wherein the nitrogen is bound to at least one additional alkyl group. The term "dialkylamino" includes groups wherein the nitrogen atom is bound to at least two additional alkyl groups. The term "arylamino" and "diarylamino" include groups wherein the nitrogen is bound to at least one or two aryl groups, respectively. The term "alkylarylamino," 15 "alkylaminoaryl" or "arylaminoalkyl" refers to an amino group which is bound to at least one alkyl group and at least one aryl group. The term "alkaminoalkyl" refers to an alkyl, alkenyl, or alkynyl group bound to a nitrogen atom which is also bound to an alkyl group.

The term "amide" or "aminocarboxy" includes compounds or moieties which contain a nitrogen atom which is bound to the carbon of a carbonyl or a thiocarbonyl group. The term 20 includes "alkaminocarboxy" groups which include alkyl, alkenyl, or alkynyl groups bound to an amino group bound to a carboxy group. It includes arylaminocarboxy groups which include aryl or heteroaryl moieties bound to an amino group which is bound to the carbon of a carbonyl or thiocarbonyl group. The terms "alkylaminocarboxy," "alkenylaminocarboxy," "alkynylaminocarboxy," and "arylamino-carboxy" include moieties wherein alkyl, alkenyl, 25 alkynyl and aryl moieties, respectively, are bound to a nitrogen atom which is in turn bound to the carbon of a carbonyl group.

The term "carbonyl" or "carboxy" includes compounds and moieties which contain a carbon connected with a double bond to an oxygen atom. Examples of moieties which contain a carbonyl include aldehydes, ketones, carboxylic acids, amides, esters, anhydrides, etc.

30 The term "thiocarbonyl" or "thiocarboxy" includes compounds and moieties which contain a carbon connected with a double bond to a sulfur atom.

The term "ether" includes compounds or moieties which contain an oxygen bonded to two different carbon atoms or heteroatoms. For example, the term includes "alkoxyalkyl" which refers to an alkyl, alkenyl, or alkynyl group covalently bonded to an oxygen atom which is covalently bonded to another alkyl group.

5 The term "ester" includes compounds and moieties which contain a carbon or a heteroatom bound to an oxygen atom which is bonded to the carbon of a carbonyl group. The term "ester" includes alkoxycarboxy groups such as methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, butoxycarbonyl, pentoxycarbonyl, etc. The alkyl, alkenyl, or alkynyl groups are as defined above.

10 The term "thioether" includes compounds and moieties which contain a sulfur atom bonded to two different carbon or hetero atoms. Examples of thioethers include, but are not limited to alkthioalkyls, alkthioalkenyls, and alkthioalkynyls. The term "alkthioalkyls" include compounds with an alkyl, alkenyl, or alkynyl group bonded to a sulfur atom which is bonded to an alkyl group. Similarly, the term "alkthioalkenyls" and alkthioalkynyls" refer to compounds
15 or moieties wherein an alkyl, alkenyl, or alkynyl group is bonded to a sulfur atom which is covalently bonded to an alkynyl group.

The term "hydroxy" or "hydroxyl" includes groups with an $-OH$ or $-O^{\cdot}$.

The term "halogen" includes fluorine, bromine, chlorine, iodine, etc. The term "perhalogenated" generally refers to a moiety wherein all hydrogens are replaced by halogen
20 atoms.

The terms "polycyclyl" or "polycyclic radical" refer to two or more cyclic rings (e.g., cycloalkyls, cycloalkenyls, cycloalkynyls, aryls and/or heterocyclyls) in which two or more carbons are common to two adjoining rings. Rings that are joined through non-adjacent atoms are termed "bridged" rings. Each of the rings of the polycycle can be substituted with such
25 substituents as described above, as for example, halogen, hydroxyl, alkylcarbonyloxy, arylcarbonyloxy, alkoxycarbonyloxy, aryloxycarbonyloxy, carboxylate, alkylcarbonyl, alkoxycarbonyl, alkylaminocarbonyl, aralkylaminocarbonyl, alkenylaminocarbonyl, alkylcarbonyl, arylcarbonyl, aralkylcarbonyl, alkenylcarbonyl, aminocarbonyl, alkylthiocarbonyl, alkoxyl, phosphate, phosphonato, phosphinato, cyano, amino (including
30 alkylamino, dialkylamino, arylamino, diarylamino, and alkylaryl amino), acylamino (including alkylcarbonylamino, arylcarbonylamino, carbamoyl and ureido), amidino, imino, sulfhydryl,

alkylthio, arylthio, thiocarboxylate, sulfates, alkylsulfinyl, sulfonato, sulfamoyl, sulfonamido, nitro, trifluoromethyl, cyano, azido, heterocyclyl, alkyl, alkylaryl, or an aromatic or heteroaromatic moiety.

The term "heteroatom" includes atoms of any element other than carbon or hydrogen.

5 Examples of heteroatoms include nitrogen, oxygen, sulfur and phosphorus.

It will be noted that the structure of some of the compounds of this invention includes asymmetric carbon atoms. It is to be understood accordingly that the isomers arising from such asymmetry (e.g., all enantiomers and diastereomers) are included within the scope of this invention, unless indicated otherwise. Such isomers can be obtained in substantially pure form
10 by classical separation techniques and by stereochemically controlled synthesis. Furthermore, the structures and other compounds and moieties discussed in this application also include all tautomers thereof.

Prodrugs are compounds which are converted *in vivo* to active forms (see, e.g., R.B. Silverman, 1992, "The Organic Chemistry of Drug Design and Drug Action", Academic Press,
15 Chp. 8). Prodrugs can be used to alter the biodistribution (e.g., to allow compounds which would not typically enter the reactive site of the protease) or the pharmacokinetics for a particular compound. For example, a hydroxyl group, can be esterified, e.g., with a carboxylic acid group to yield an ester. When the ester is administered to a subject, the ester is cleaved, enzymatically or non-enzymatically, reductively or hydrolytically, to reveal the hydroxyl
20 group.

The term "prodrug moiety" includes moieties which can be metabolized *in vivo* to a hydroxyl group and moieties which may advantageously remain esterified *in vivo*. Preferably, the prodrugs moieties are metabolized *in vivo* by esterases or by other mechanisms to hydroxyl groups or other advantageous groups. Examples of prodrugs and their uses are well known in
25 the art (See, e.g., Berge et al. (1977) "Pharmaceutical Salts", *J. Pharm. Sci.* 66:1-19). The prodrugs can be prepared *in situ* during the final isolation and purification of the compounds, or by separately reacting the purified compound in its free acid form or hydroxyl with a suitable esterifying agent. Hydroxyl groups can be converted into esters *via* treatment with a carboxylic acid. Examples of prodrug moieties include substituted and unsubstituted, branch or
30 unbranched lower alkyl ester moieties, (e.g., propionic acid esters), lower alkenyl esters, di-lower alkyl-amino lower-alkyl esters (e.g., dimethylaminoethyl ester), acylamino lower alkyl esters (e.g., acetyloxymethyl ester), acyloxy lower alkyl esters (e.g., pivaloyloxymethyl ester),

aryl esters (phenyl ester), aryl-lower alkyl esters (e.g., benzyl ester), substituted (e.g., with methyl, halo, or methoxy substituents) aryl and aryl-lower alkyl esters, amides, lower-alkyl amides, di-lower alkyl amides, and hydroxy amides. Preferred prodrug moieties are propionic acid esters and acyl esters.

5 The invention also features a method for treating a tetracycline compound responsive state in a subject, by administering to the subject a 7-substituted tetracycline compound of the invention, e.g., a compound of formula I. Preferably, an effective amount of the tetracycline compound is administered. Examples of 7-substituted tetracycline compounds of the invention include 7-phenyl sancycline, 7, 9 diphenyl sancycline, 7-(2-fluorophenyl) sancycline, 7-(2-
10 chlorophenyl) sancycline, 7-(2-bromophenyl) sancycline, 7-(2-iodophenyl) sancycline, 7-(3-fluorophenyl) sancycline, 7-(3-chlorophenyl) sancycline, 7-(3-bromophenyl) sancycline, 7-(3-iodophenyl) sancycline, 7-(4-fluorophenyl) sancycline, 7-(4-chlorophenyl) sancycline, 7-(4-bromophenyl) sancycline, 7-(4-iodophenyl) sancycline, 7-(4-trichloromethylphenyl)
sancycline, 7-(4-trifluoromethylphenyl) sancycline, 7-(4-tribromomethylphenyl) sancycline, or
15 7-(4-triiodomethylphenyl) sancycline.

 The language "tetracycline compound responsive state" includes states which can be treated, prevented, or otherwise ameliorated by the administration of a tetracycline compound of the invention. Tetracycline compound responsive states include bacterial infections (including those which are resistant to other tetracycline compounds), cancer, diabetes, and
20 other states for which tetracycline compounds have been found to be active (see, for example, U.S. Patent Nos. 5,789,395; 5,834,450; and 5,532,227). Compounds of the invention can be used to prevent or control important mammalian and veterinary diseases such as diarrhea, urinary tract infections, infections of skin and skin structure, ear, nose and throat infections, wound infection, mastitis and the like. In addition, methods for treating neoplasms using
25 tetracycline compounds of the invention are also included (van der Bozert et al., *Cancer Res.*, 48:6686-6690 (1988)).

 Bacterial infections may be caused by a wide variety of gram positive and gram negative bacteria. The compounds of the invention are useful as antibiotics against organisms which are resistant to other tetracycline compounds. The antibiotic activity of the tetracycline
30 compounds of the invention may be determined using the method discussed in Example 2, or by using the in vitro standard broth dilution method described in Waitz, J.A., *National*

Commission for Clinical Laboratory Standards, Document M7-A2, vol. 10, no. 8, pp. 13-20, 2nd edition, Villanova, PA (1990).

The tetracycline compounds may also be used to treat infections traditionally treated with tetracycline compounds such as, for example, rickettsiae; a number of gram-positive and gram-negative bacteria; and the agents responsible for lymphogranuloma venereum, inclusion conjunctivitis, psittacosis. The tetracycline compounds may be used to treat infections of, e.g., *K. pneumoniae*, *Salmonella*, *E. hirae*, *A. baumannii*, *B. catarrhalis*, *H. influenzae*, *P. aeruginosa*, *E. faecium*, *E. coli*, *S. aureus* or *E. faecalis*. In one embodiment, the tetracycline compound is used to treat a bacterial infection that is resistant to other tetracycline antibiotic compounds. The tetracycline compound of the invention may be administered with a pharmaceutically acceptable carrier.

The language "effective amount" of the compound is that amount necessary or sufficient to treat or prevent a tetracycline compound responsive state. The effective amount can vary depending on such factors as the size and weight of the subject, the type of illness, or the particular tetracycline compound. For example, the choice of the tetracycline compound can affect what constitutes an "effective amount". One of ordinary skill in the art would be able to study the aforementioned factors and make the determination regarding the effective amount of the tetracycline compound without undue experimentation.

The invention also pertains to methods of treatment against microorganism infections and associated diseases. The methods include administration of an effective amount of one or more tetracycline compounds to a subject. The subject can be either a plant or, advantageously, an animal, e.g., a mammal, e.g., a human.

In the therapeutic methods of the invention, one or more tetracycline compounds of the invention may be administered alone to a subject, or more typically a compound of the invention will be administered as part of a pharmaceutical composition in mixture with conventional excipient, i.e., pharmaceutically acceptable organic or inorganic carrier substances suitable for parenteral, oral or other desired administration and which do not deleteriously react with the active compounds and are not deleterious to the recipient thereof.

In one embodiment, the pharmaceutical composition comprises a 7-substituted tetracycline compound of the invention, e.g., of formula I. In a further embodiment, the 7-substituted tetracycline compound is 7-phenyl sancycline, 7, 9 diphenyl sancycline, 7-(2-

fluorophenyl) sancycline, 7-(2-chlorophenyl) sancycline, 7-(2-bromophenyl) sancycline, 7-(2-iodophenyl) sancycline, 7-(3-fluorophenyl) sancycline, 7-(3-chlorophenyl) sancycline, 7-(3-bromophenyl) sancycline, 7-(3-iodophenyl) sancycline, 7-(4-fluorophenyl) sancycline, 7-(4-chlorophenyl) sancycline, 7-(4-bromophenyl) sancycline, 7-(4-iodophenyl) sancycline, 7-(4-trichloromethylphenyl) sancycline, 7-(4-trifluoromethylphenyl) sancycline, 7-(4-tribromomethylphenyl) sancycline, or 7-(4-triiodomethylphenyl) sancycline.

The language "pharmaceutically acceptable carrier" includes substances capable of being coadministered with the tetracycline compound(s), and which allow both to perform their intended function, e.g., treat or prevent a tetracycline compound responsive state. Suitable
10 pharmaceutically acceptable carriers include but are not limited to water, salt solutions, alcohol, vegetable oils, polyethylene glycols, gelatin, lactose, amylose, magnesium stearate, talc, silicic acid, viscous paraffin, perfume oil, fatty acid monoglycerides and diglycerides, petroethral fatty acid esters, hydroxymethyl-cellulose, polyvinylpyrrolidone, etc. The pharmaceutical preparations can be sterilized and if desired mixed with auxiliary agents, e.g.,
15 lubricants, preservatives, stabilizers, wetting agents, emulsifiers, salts for influencing osmotic pressure, buffers, colorings, flavorings and/or aromatic substances and the like which do not deleteriously react with the active compounds of the invention.

The tetracycline compounds of the invention that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be
20 used to prepare pharmaceutically acceptable acid addition salts of the tetracycline compounds of the invention that are basic in nature are those that form non-toxic acid addition salts, i.e., salts containing pharmaceutically acceptable anions, such as the hydrochloride, hydrobromide, hydroiodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, acid citrate, tartrate, pantothenate, bitartrate, ascorbate, succinate, maleate,
25 gentisinate, fumarate, gluconate, glucaronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, p-toluenesulfonate and palmoate [i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)] salts. Although such salts must be pharmaceutically acceptable for administration to a subject, e.g., a mammal, it is often desirable in practice to initially isolate a tetracycline compound of the invention from the
30 reaction mixture as a pharmaceutically unacceptable salt and then simply convert the latter back to the free base compound by treatment with an alkaline reagent and subsequently convert the latter free base to a pharmaceutically acceptable acid addition salt. The acid addition salts

of the base compounds of this invention are readily prepared by treating the base compound with a substantially equivalent amount of the chosen mineral or organic acid in an aqueous solvent medium or in a suitable organic solvent, such as methanol or ethanol. Upon careful evaporation of the solvent, the desired solid salt is readily obtained. The preparation of other
5 tetracycline compounds of the invention not specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

The preparation of other tetracycline compounds of the invention not specifically described in the foregoing experimental section can be accomplished using combinations of the
10 reactions described above that will be apparent to those skilled in the art.

The tetracycline compounds of the invention that are acidic in nature are capable of forming a wide variety of base salts. The chemical bases that may be used as reagents to prepare pharmaceutically acceptable base salts of those tetracycline compounds of the invention that are acidic in nature are those that form non-toxic base salts with such
15 compounds. Such non-toxic base salts include, but are not limited to those derived from such pharmaceutically acceptable cations such as alkali metal cations (e.g., potassium and sodium) and alkaline earth metal cations (e.g., calcium and magnesium), ammonium or water-soluble amine addition salts such as N-methylglucamine-(meglumine), and the lower alkanolammonium and other base salts of pharmaceutically acceptable organic amines. The
20 pharmaceutically acceptable base addition salts of tetracycline compounds of the invention that are acidic in nature may be formed with pharmaceutically acceptable cations by conventional methods. Thus, these salts may be readily prepared by treating the tetracycline compound of the invention with an aqueous solution of the desired pharmaceutically acceptable cation and evaporating the resulting solution to dryness, preferably under reduced pressure. Alternatively,
25 a lower alkyl alcohol solution of the tetracycline compound of the invention may be mixed with an alkoxide of the desired metal and the solution subsequently evaporated to dryness.

The preparation of other tetracycline compounds of the invention not specifically described in the foregoing experimental section can be accomplished using combinations of the reactions described above that will be apparent to those skilled in the art.

30 The tetracycline compounds of the invention and pharmaceutically acceptable salts thereof can be administered via either the oral, parenteral or topical routes. In general, these compounds are most desirably administered in effective dosages, depending upon the weight

and condition of the subject being treated and the particular route of administration chosen. Variations may occur depending upon the species of the subject being treated and its individual response to said medicament, as well as on the type of pharmaceutical formulation chosen and the time period and interval at which such administration is carried out.

5 The pharmaceutical compositions of the invention may be administered alone or in combination with other known compositions for treating tetracycline responsive states in a mammal. Preferred mammals include pets (e.g., cats, dogs, ferrets, etc.), farm animals (cows, sheep, pigs, horses, goats, etc.), lab animals (rats, mice, monkeys, etc.), and primates (chimpanzees, humans, gorillas). The language "in combination with" a known composition is
10 intended to include simultaneous administration of the composition of the invention and the known composition, administration of the composition of the invention first, followed by the known composition and administration of the known composition first, followed by the composition of the invention. Any of the therapeutically composition known in the art for treating tetracycline responsive states can be used in the methods of the invention.

15 The compounds of the invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by any of the routes previously mentioned, and the administration may be carried out in single or multiple doses. For example, the novel therapeutic agents of this invention can be administered advantageously in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable
20 inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various non-toxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the
25 therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging from about 5.0% to about 70% by weight.

 For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dicalcium phosphate and glycine may be employed along with various disintegrants such as starch (and preferably corn, potato or
30 tapioca starch), alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes.

Solid compositions of a similar type may also be employed as fillers in gelatin capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or
5 flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration (including intraperitoneal, subcutaneous, intravenous, intradermal or intramuscular injection), solutions of a therapeutic compound of the present
10 invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH greater than 8) if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intraarticular, intramuscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions
15 is readily accomplished by standard pharmaceutical techniques well known to those skilled in the art. For parenteral application, examples of suitable preparations include solutions, preferably oily or aqueous solutions as well as suspensions, emulsions, or implants, including suppositories. Therapeutic compounds may be formulated in sterile form in multiple or single dose formats such as being dispersed in a fluid carrier such as sterile physiological saline or 5%
20 saline dextrose solutions commonly used with injectables.

Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin. Examples of methods of topical administration include transdermal, buccal or sublingual application. For topical applications, therapeutic compounds can be suitably admixed in a pharmacologically inert topical carrier
25 such as a gel, an ointment, a lotion or a cream. Such topical carriers include water, glycerol, alcohol, propylene glycol, fatty alcohols, triglycerides, fatty acid esters, or mineral oils. Other possible topical carriers are liquid petrolatum, isopropylpalmitate, polyethylene glycol, ethanol 95%, polyoxyethylene monolaurate 5% in water, sodium lauryl sulfate 5% in water, and the like. In addition, materials such as anti-oxidants, humectants, viscosity stabilizers and the like
30 also may be added if desired.

For enteral application, particularly suitable are tablets, dragees or capsules having talc and/or carbohydrate carrier binder or the like, the carrier preferably being lactose and/or corn

starch and/or potato starch. A syrup, elixir or the like can be used wherein a sweetened vehicle is employed. Sustained release compositions can be formulated including those wherein the active component is protected with differentially degradable coatings, e.g., by microencapsulation, multiple coatings, etc.

5 In addition to treatment of human subjects, the therapeutic methods of the invention also will have significant veterinary applications, e.g. for treatment of livestock such as cattle, sheep, goats, cows, swine and the like; poultry such as chickens, ducks, geese, turkeys and the like; horses; and pets such as dogs and cats. Also, the compounds of the invention may be used to treat non-animal subjects, such as plants.

10 It will be appreciated that the actual preferred amounts of active compounds used in a given therapy will vary according to the specific compound being utilized, the particular compositions formulated, the mode of application, the particular site of administration, etc. Optimal administration rates for a given protocol of administration can be readily ascertained by those skilled in the art using conventional dosage determination tests conducted with regard
15 to the foregoing guidelines.

In general, compounds of the invention for treatment can be administered to a subject in dosages used in prior tetracycline therapies. See, for example, the *Physicians' Desk Reference*. For example, a suitable effective dose of one or more compounds of the invention will be in the range of from 0.01 to 100 milligrams per kilogram of body weight of recipient per day,
20 preferably in the range of from 0.1 to 50 milligrams per kilogram body weight of recipient per day, more preferably in the range of 1 to 20 milligrams per kilogram body weight of recipient per day. The desired dose is suitably administered once daily, or several sub-doses, e.g. 2 to 5 sub-doses, are administered at appropriate intervals through the day, or other appropriate schedule.

25 It will also be understood that normal, conventionally known precautions will be taken regarding the administration of tetracyclines generally to ensure their efficacy under normal use circumstances. Especially when employed for therapeutic treatment of humans and animals *in vivo*, the practitioner should take all sensible precautions to avoid conventionally known contradictions and toxic effects. Thus, the conventionally recognized adverse reactions of
30 gastrointestinal distress and inflammations, the renal toxicity, hypersensitivity reactions, changes in blood, and impairment of absorption through aluminum, calcium, and magnesium ions should be duly considered in the conventional manner.

Furthermore, the invention also pertains to the use of a tetracycline compound of formula I, for the preparation of a medicament. The medicament may include a pharmaceutically acceptable carrier and the tetracycline compound is an effective amount, e.g., an effective amount to treat a tetracycline responsive state.

5 In yet another embodiment, the invention also pertains to the use of a tetracycline compound of formula I to treat a tetracycline responsive state, e.g., in a subject, e.g., a mammal, e.g., a human.

Compounds of the invention may be made as described below, with modifications to the procedure below within the skill of those of ordinary skill in the art.

10 **EXAMPLE 1: Synthesis of the 7- Substituted Tetracycline Compounds**

Preparation of 7-iodosancycline and 7, 9-diiodosancycline

Reaction of sancycline with N-iodosuccinimide:

Five grams of sancycline was dissolved in 85 mL of concentrated sulfuric acid that was cooled to 0°C (on ice). N-iodosuccinimide (NIS) was added to the reaction in 300 mg portions every 15 minutes and reacted for 5 hours. The reaction is removed from the ice bath. The mixture, analyzed by HPLC or TLC, show the product of D-ring iodotetracyclines. After the reaction was complete, the sulfuric acid was dripped slowly 1 L of ice water and extracted 7 times with 300 mL of n-butanol. The solvent was removed *in vacuo* to produce a mixture of three products. The 7-iodo regioisomer, 9-regioisomer and 7,9-diiodosancycline derivative of sancycline were purified by preparative HPLC chromatography or by methods known in the art.

Rt: Hypersil C18 BDS Column: 7 and 9 isomer mixture: in the ratio 40:60 for position 9/position 7

25 7-iodosancycline: Rt 14.45 min MS (M+H, formic acid solvent): 541.1
¹H NMR (Methanol d₄-300 MHz) δ 7.89 (d, J=8.86 Hz, 1H), 6.67 (d, 8.87 Hz, 1H), 3.78 (s, 1H), 3.03 (s, 2H), 2.84 (s, 6H), 2.46 (m, 2H), 1.63 (m, 4H), 0.95 (m, 2H)

30 9-iodosancycline: Rt 14.1 min: MS (M+H, formic acid solvent): 541.1
¹H NMR (Methanol d₄-300 MHz) δ 7.87 (d, J=8.86 Hz, 1H), 6.64 (d, 8.87 Hz, 1H), 3.78 (s, 1H), 3.03 (s, 2H), 2.84 (s, 6H), 2.46 (m, 2H), 1.63 (m, 4H), 0.95 (m, 2H)

35

7,9-diiodo sancycline: Rt 21.2 min

MS (M+H, formic acid solvent): 667.3

¹H NMR (Methanol d₄-300 MHz) δ 8.35, 3.78 (s, 1H), 3.33 (s, 2H), 2.88 (s, 7H), 2.41 (m, 2H), 1.41 (m, 5H).

5

7-phenylsancycline

7-iodosancycline, 150 mg (0.28 mM), Pd(OAc)₂ and 10 mL of MeOH are added to a flask with a stir bar and the system degassed 3x using argon. Na₂CO₃ (87 mg, 0.8mM) dissolved in water and argon degassed is added via syringe is added along with phenylboronic acid (68 mg, 0.55 mM) in MeOH that was also degassed. The reaction was followed by HPLC for 2 hours and cooled to room temperature. The solution was filtered, and dried to produce a crude mixture. The solid was dissolved in dimethylformamide and injected onto a preparative HPLC system using C18 reverse-phase silica. The fraction at 36-38 minutes was isolated, and the solvent removed *in vacuo* to yield the product plus salts. The salts were removed before extraction into 50:25:25 water:butanol:ethyl acetate and dried *in vacuo*. This solid was dissolved in MeOH and the HCl salt made by bubbling in HCl gas. The solvent was removed to produce the product in 42% yield as a yellow solid.

7-phenyl sancycline: Rt 21.6 min: MS (M+H, formic acid solvent): 491.3
¹H NMR (Methanol d₄-300 MHz) δ 7.87 (d, J=8.86 Hz, 1H), 7.38 (m, 5H), 6.64 (d, 8.87 Hz, 1H), 4.00 (s, 1H), 3.84 (s, 2H), 3.01 (s, 6H), 2.46 (m, 2H), 1.63 (m, 4H), 0.95 (m, 2H)

7-(4-chlorophenyl)sancycline

7-iodosancycline, 500 mg (0.91 mM), Pd(OAc)₂ 21 mg, and 20 mL of MeOH are added to a flask with a stir bar and the system degassed 3x using argon. Na₂CO₃ (293mg, 2.8mM) dissolved in water and argon degassed, is added via syringe is added along with 4-Cl-phenylboronic acid (289 mg, 1.85 mM) in MeOH that was also degassed. The reaction was followed by HPLC for 45 minutes and cooled to room temperature. The solution was filtered, and dried to produce a crude mixture. The solid was dissolved in dimethylformamide and injected onto a preparative HPLC system using C18 reverse-phase silica. The fraction at 39 minutes was isolated, and the solvent removed *in vacuo* to yield the product plus salts. The salts were removed by extraction into 50:25:25 water:butanol:ethyl acetate, and dried *in vacuo*. This solid was dissolved in MeOH and the HCl salt made by bubbling in HCl gas. The solvent was removed to produce the product in 57% yield as a yellow solid.

7-(4-chlorophenyl)sancycline: Rt 20.3 min: MS (M+H, formic acid solvent): 525.7
¹H NMR (Methanol d₄-300 MHz)δ 7.49-7.52 (d, J=8.54 Hz, 1H), 6.99-7.01 (d, 8.61 Hz, 1H), 4.12 (s, 1H), 3.67 (m, 1H), 3.06 (s, 6H), 2.58 (m, 2H), 1.62(m, 4H), 1.01 (m, 2H)

5

7-(4-fluorophenyl)sancycline

7-iodosancycline, 200 mg (0.3 mM), Pd(OAc)₂ 8.3 mg, and 10 mL of MeOH are added to a flask with a stir bar and the system degassed 3x using argon. Na₂CO₃ (104 mg, 1.1 mM) dissolved in water and argon degassed is added via syringe is added along with 4-F-phenylboronic acid (104 mg, 0.7 mM) in MeOH that was also degassed. The reaction was followed by HPLC for 20 minutes and cooled to room temperature. The solution was filtered, and dried to produce a crude mixture. The solid was dissolved in dimethylformamide and injected onto a preparative HPLC system using C18 reverse-phase silica. The fraction at 19-20 minutes was isolated, and the solvent removed *in vacuo* to yield the product plus salts. The salts were removed by extraction into 50:25:25 water:butanol:ethyl acetate and dried *in vacuo*. This solid was dissolved in MeOH and the HCl salt made by bubbling in HCl gas. The solvent was removed to produce the product in 47% yield as a yellow solid.

7-(4-fluorophenyl)sancycline: Rt 19.5 min: MS (M+H, formic acid solvent): 509.4
¹H NMR (Methanol d₄-300 MHz)δ 6.92-6.95 (d, 1H), 7.45-7.48 (d, 1H), 7.15-7.35 (m, 4H), 4.05 (s, 1H), 3.62 (m, 1H), 3.08 (s, 6H), 2.55 (m, 2H), 1.65(m, 4H), 1.00 (m, 2H)

20

Example 2: *In vitro* Minimum Inhibitory Concentration (MIC) Assay

The following assay is used to determine the efficacy of tetracycline compounds against common bacteria. 2 mg of each compound is dissolved in 100 µl of DMSO. The solution is then added to cation-adjusted Mueller Hinton broth (CAMHB), which results in a final compound concentration of 200 µg per ml. The tetracycline compound solutions are diluted to 50 µL volumes, with a test compound concentration of .098 µg/ml. Optical density (OD) determinations are made from fresh log-phase broth cultures of the test strains. Dilutions are made to achieve a final cell density of 1x10⁶ CFU/ml. At OD=1, cell densities for different genera should be approximately:

30

35

<i>E. coli</i>	1x10 ⁹ CFU/ml
<i>S. aureus</i>	5x10 ⁸ CFU/ml
<i>Enterococcus sp.</i>	2.5x10 ⁹ CFU/ml

50 μ l of the cell suspensions are added to each well of microtiter plates. The final cell density should be approximately 5×10^5 CFU/ml. These plates are incubated at 35°C in an ambient air incubator for approximately 18 hr. The plates are read with a microplate reader and are visually inspected when necessary. The MIC is defined as the lowest concentration of the tetracycline compound that inhibits growth. Compounds of the invention indicate good inhibition of growth.

EQUIVALENTS


Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, numerous equivalents to the specific procedures described herein.

10 Such equivalents are considered to be within the scope of the present invention and are covered by the following claims. The contents of all references, issued patents, and published patent applications cited throughout this application are hereby incorporated by reference. The appropriate components, processes, and methods of those patents, applications and other documents may be selected for the present invention and embodiments thereof.

CLAIMS

What is claimed is:

1. A 7-substituted tetracycline compound of the formula:



5

wherein:

R^4 and $R^{4'}$ are each alkyl;

R^5 is hydrogen, hydroxyl, or a prodrug moiety;

10

R^6 and $R^{6'}$ are each independently hydrogen, hydroxyl, alkyl, or taken together, alkenyl;

R^7 is halo substituted or unsubstituted phenyl;

and pharmaceutically acceptable salts thereof.

15

2. The compound of claim 1, wherein R^5 , R^6 and $R^{6'}$ are each hydrogen and R^4 and $R^{4'}$ are each methyl.

3. The compound of claim 1, wherein R^7 is unsubstituted phenyl.

4. The compound of claim 3, wherein said compound is selected from the group consisting of 7-phenylsancycline and 7, 9 diphenyl sancycline.

5. The compound of claim 1, wherein R^7 is 2-substituted phenyl.

20


6. The compound of claim 5, wherein said compound is selected from the group consisting of 7-(2-fluorophenyl) sancycline, 7-(2-chlorophenyl) sancycline, 7-(2-bromophenyl) sancycline, and 7-(2-iodophenyl) sancycline.

7. The compound of claim 1, wherein R^7 is 3-substituted phenyl.

25

8. The compound of claim 7, wherein said compound is selected from the group consisting of 7-(3-fluorophenyl) sancycline, 7-(3-chlorophenyl) sancycline, 7-(3-bromophenyl) sancycline, and 7-(3-iodophenyl) sancycline.

9. The compound of claim 1, wherein R^7 is 4-substituted phenyl.
10. The compound of claim 9, wherein said compound is selected from the group consisting of 7-(4-fluorophenyl) sancycline, 7-(4-chlorophenyl) sancycline, 7-(4-bromophenyl) sancycline, 7-(4-iodophenyl) sancycline, 7-(4-trichloromethylphenyl) sancycline, 7-(4-trifluoromethylphenyl) sancycline, 7-(4-tribromomethylphenyl) sancycline, and 7-(4-triiodomethylphenyl) sancycline.
11. A method for treating a tetracycline responsive state in a mammal, comprising administering to said mammal a 7-substituted tetracycline compound of formula (I):



10 wherein:

R^4 and $R^{4'}$ are each alkyl;

R^5 is hydrogen, hydroxyl, or a prodrug moiety;

R^6 and $R^{6'}$ are each independently hydrogen, hydroxyl, alkyl, or taken together, alkenyl;

15 R^7 is halo substituted or unsubstituted phenyl; and pharmaceutically acceptable salts thereof, such that the tetracycline responsive state is treated.

12. The method of claim 11, wherein R^5 , R^6 and $R^{6'}$ are each hydrogen and R^4 and $R^{4'}$ are each methyl.
13. The method of claim 11, wherein R^7 is unsubstituted phenyl.
14. The method of claim 13, wherein said compound is selected from the group consisting of 7-phenylsancycline and 7, 9 diphenyl sancycline.
- 20 15. The method of claim 1, wherein R^7 is 2-substituted phenyl.
16. The method of claim 15, wherein said compound is selected from the group consisting of 7-(2-fluorophenyl) sancycline, 7-(2-chlorophenyl) sancycline, 7-(2-bromophenyl) sancycline, and 7-(2-iodophenyl) sancycline.
- 25 17. The method of claim 11, wherein R^7 is 3-substituted phenyl.

18. The method of claim 17, wherein said compound is selected from the group consisting of 7-(3-fluorophenyl) sancycline, 7-(3-chlorophenyl) sancycline, 7-(3-bromophenyl) sancycline, and 7-(3-iodophenyl) sancycline.
19. The method of claim 11, wherein R⁷ is 4-substituted phenyl.
- 5 20. The method of claim 19, wherein said compound is selected from the group consisting of 7-(4-fluorophenyl) sancycline, 7-(4-chlorophenyl) sancycline, 7-(4-bromophenyl) sancycline, 7-(4-iodophenyl) sancycline, 7-(4-trichloromethylphenyl) sancycline, 7-(4-trifluoromethylphenyl) sancycline, 7-(4-tribromomethylphenyl) sancycline, and 7-(4-triiodomethylphenyl) sancycline.
- 10 21. The method of claim 11, wherein said tetracycline responsive state is a bacterial infection.
22. The method of claim 21, wherein said bacterial infection is associated with *E. coli*.
23. The method of claim 21, wherein said bacterial infection is associated with *S. aureus*.
24. The method of claim 21, wherein said bacterial infection is associated with *E. faecalis*.
- 15 25. The method of claim 21, wherein said bacterial infection is resistant to other tetracycline antibiotics.
26. The method of claim 11, wherein said compound is administered with a pharmaceutically acceptable carrier.
27. A pharmaceutical composition comprising a therapeutically effective amount of a
20 compound of claim 1 and a pharmaceutically acceptable carrier.
28. The pharmaceutical composition of claim 27, wherein said compound is selected from the group consisting of 7-phenyl sancycline, 7, 9 diphenyl sancycline, 7-(2-fluorophenyl) sancycline, 7-(2-chlorophenyl) sancycline, 7-(2-bromophenyl) sancycline, 7-(2-iodophenyl) sancycline, 7-(3-fluorophenyl) sancycline, 7-(3-chlorophenyl) sancycline, 7-(3-bromophenyl) sancycline, 7-(3-iodophenyl) sancycline,
25 7-(4-fluorophenyl) sancycline, 7-(4-chlorophenyl) sancycline, 7-(4-bromophenyl) sancycline, 7-(4-iodophenyl) sancycline, 7-(4-trichloromethylphenyl) sancycline, 7-(4-trifluoromethylphenyl) sancycline, 7-(4-tribromomethylphenyl) sancycline, and 7-(4-triiodomethylphenyl) sancycline.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/16632

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C237/26 A61K31/165 A61P31/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	KOZA ET AL: "Synthesis of 7-Substituted Tetracyclin Derivatives" ORGANIC LETTERS, ACS, WASHINGTON, DC, US, vol. 2, no. 6, 1 March 2000 (2000-03-01), pages 815-817, XP000943852 ISSN: 1523-7060 the whole document	1-4, 9, 10

☐ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

A document member of the same patent family

Date of the actual completion of the international search

11 January 2001

Date of mailing of the international search report

18/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Rufet, J